

**DAHLGREN DIVISION
NAVAL SURFACE WARFARE CENTER**

Dahlgren, Virginia 22448-5100




NSWCDD/TR-94/77

**IMPROVED METHOD FOR CALCULATING EXACT
GEODETIC LATITUDE AND ALTITUDE**

BY ISAAC SOFAIR

STRATEGIC AND SPACE SYSTEMS DEPARTMENT

APRIL 1995



Approved for public release; distribution is unlimited.

19950607 053

DTIC QUALITY INSPECTED 8

FOREWORD

The work described in this report was performed in the Fire Control Formulation Branch (K41), Submarine Launched Ballistic Missile (SLBM) Research and Analysis Division, Strategic and Space Systems Department, and was authorized under Strategic Systems Program Office Task Assignment 36401. This work was necessitated by the need to formulate an exact method of computing geodetic latitude and altitude whose chief attribute is the absence of singularities in the results.

This report is a refined version of NSWC TR 85-85, dated April 1985 and revised May 1986. NSWC TR 85-85 should be discarded.

This report has been reviewed and approved by Davis Owen; Johnny Boyles; Carol Rose, Head, Fire Control Formulation Branch; and Dr. David Lando, Head, SLBM Research and Analysis Division.

Approved by:



R. L. SCHMIDT, Head
Strategic and Space Systems Department

INTRODUCTION

A simple and efficient model is presented for calculating the exact geodetic latitude and altitude of an arbitrary point in space, given the coordinates of that point. The model assumes the Earth to be an oblate spheroid; i.e., an ellipsoid of revolution whose semimajor axis 'a' is the radius of the circle described by the equatorial plane, and whose semiminor axis 'b' is a line joining its center and one of its poles. The point in question can be either inside or outside the Earth ellipsoid, excluding a small open region bounded by a prolate spheroid that is concentric with and contained within the Earth ellipsoid. The cardinal merit of this model is the absence of singularities in the results.

FORMULATION

Choose an Earth-fixed Cartesian coordinate system whose origin coincides with the center of the Earth ellipsoid. The unit vectors \hat{i} , \hat{j} , and \hat{k} coincide with the x, y, and z axes, respectively. The +z axis points in the direction of the North Pole. The +x axis is the line of intersection of the equatorial plane with the plane of zero longitude. The +y axis completes a right-handed coordinate system.

An equation of the ellipsoid in this frame is

$$\frac{r^2}{a^2} + \frac{z^2}{b^2} = 1, \text{ where } r = \sqrt{x^2 + y^2}. \quad (1)$$

Let $P(x_0, y_0, z_0)$ be the coordinates of the given point. It is desired to find the nearest point $P'(x, y, z)$ to P on the surface of the ellipsoid.

The slope of a normal to the ellipsoid at any point on its surface is given by

$$-\frac{1}{\frac{dz}{dr}} = \frac{a^2 z}{b^2 r}, \text{ where } \frac{r^2}{a^2} + \frac{z^2}{b^2} = 1. \quad (2)$$

Therefore, the slope of a normal from P is

$$\frac{z_0 - z}{r_0 - r} = \frac{a^2 z}{b^2 r}, \text{ where } r_0 = \sqrt{x_0^2 + y_0^2};$$

i.e.,

$$(r_0 - r)a^2 z = (z_0 - z)b^2 r,$$

or

$$r_0 a^2 z = r \{ (a^2 - b^2)z + b^2 z_0 \} .$$

Squaring both sides and expressing r in terms of z ,

$$a^2 b^2 r_0^2 z^2 = (b^2 - z^2) \{ (a^2 - b^2)z + b^2 z_0 \}^2 .$$

Writing the above equation in descending powers of z , the following quartic in z is obtained:

$$(a^2 - b^2)^2 z^4 + 2b^2(a^2 - b^2)z_0 z^3 + b^2 \{ a^2 r_0^2 + b^2 z_0^2 - (a^2 - b^2)^2 \} z^2 - 2b^4(a^2 - b^2)z_0 z - b^6 z_0^2 = 0 .$$

Since z is either positive or negative and $\text{sign } z = \text{sign } z_0$, the above quartic can be expressed in terms of $|z|$. The result is

$$|z|^4 + 2p|z|^3 + q|z|^2 - 2pb^2|z| - p^2b^2 = 0 , \quad (3)$$

where

$$p = \frac{b^2 |z_0|}{a^2 - b^2} = \frac{|z_0|}{e'^2} , \quad (4)$$

$$q = \frac{b^2 \{ a^2 r_0^2 + b^2 z_0^2 - (a^2 - b^2)^2 \}}{(a^2 - b^2)^2} = p^2 - b^2 + \frac{r_0^2}{e^2 e'^2} , \quad (5)$$

where

$$e^2 = 1 - \frac{b^2}{a^2} , \quad e'^2 = \frac{a^2}{b^2} - 1 .$$

Two of three powerful theorems in the Theory of Equations will now be employed to expose the nature of the roots of Equation (3). The third theorem will be needed later on. These theorems are found in the reference¹.

I. An equation $f(x) = 0$ cannot have more positive roots than there are changes of sign in $f(x)$, and cannot have more negative roots than there are changes of sign in $f(-x)$.

II. Every equation that is of an even degree and has its last term negative has at least two real roots, one positive and one negative.

III. Every equation of an odd degree has at least one real root whose sign is opposite to that of its last term.

By observing Equations (3), (4), and (5), the following conclusions are deduced:

¹ Hall, H.S. and Knight, S.R., Higher Algebra, Fourth Edition, MacMillan and Co., London, 1927.

(i) Since the last term of Equation (3) is negative, there are at least two real roots of opposite sign (using II).

(ii) Since there is only one change of sign in Equation (3), there is at most one positive root (using I).

From (i) and (ii), it is seen immediately that Equation (3) has exactly one positive root, and this is the root that is sought.

The solution of Equation (3) is effected by a standard method known as Ferrari's method¹. In the equation

$$|z|^4 + 2p|z|^3 + q|z|^2 - 2pb^2|z| - p^2b^2 = 0 ,$$

add to each side $(c|z|+d)^2$, the quantities c and d being determined so as to make the left-hand side a perfect square; then

$$|z|^4 + 2p|z|^3 + (q + c^2)|z|^2 + 2(cd - pb^2)|z| + d^2 - p^2b^2 = (c|z| + d)^2 .$$

Suppose that the left side of the equation is equal to $(|z|^2 + p|z| + t)^2$, then comparing the coefficients,

$$p^2 + 2t = q + c^2, \quad pt = cd - pb^2, \quad t^2 = d^2 - p^2b^2 . \quad (6)$$

Eliminating c and d from these equations,

$$p^2(t + b^2)^2 = (2t + p^2 - q)(t^2 + p^2b^2) ,$$

or

$$2t^3 - qt^2 - p^2b^2(q - p^2 + b^2) = 0 ,$$

or

$$2t^3 - qt^2 - \frac{p^2b^2r_0^2}{e^2e'^2} = 0, \text{ using Equation (5)}$$

or

$$2t^3 - qt^2 - \frac{a^2r_0^2z_0^2}{e'^8} = 0 . \quad (7)$$

Applying Theorems I and III to Equation (7), we see that Equation (7) has exactly one positive root. The solution of Equation (7) is accomplished by a standard method known as Cardan's Solution¹. Eliminating the t^2 term in Equation (7) by making the substitution

$$t = t' + \frac{q}{6} ;$$

i.e.,

$$2\left(t' + \frac{q}{6}\right)^3 - q\left(t' + \frac{q}{6}\right)^2 - \frac{a^2 r_0^2 z_0^2}{e'^8} = 0 ,$$

then

$$t'^3 + Ft' + H = 0 , \quad (8)$$

where

$$F = -\frac{q^2}{12}, H = -\frac{q^3}{108} - \frac{a^2 r_0^2 z_0^2}{2e'^8} .$$

The solution of Equation (8) is

$$t' = \left(\sqrt{\frac{H^2}{4} + \frac{F^3}{27}} - \frac{H}{2} \right)^{\frac{1}{3}} - \left(\sqrt{\frac{H^2}{4} + \frac{F^3}{27}} + \frac{H}{2} \right)^{\frac{1}{3}} . \quad (9)$$

Equation (9) is valid provided that

$$\frac{H^2}{4} + \frac{F^3}{27} \geq 0 .$$

We have

$$H = -\frac{q^3}{108} - \frac{a^2 r_0^2 z_0^2}{2e'^8} = -\frac{2}{6^3}(q^3 + 2P) ,$$

where

$$P = \frac{27a^2 r_0^2 z_0^2}{e'^8} .$$

$$\frac{H^2}{4} + \frac{F^3}{27} = \frac{1}{6^6}(q^3 + 2P)^2 - \frac{q^6}{6^6} = \frac{4}{6^6}P(P + q^3) .$$

If

$$\frac{H^2}{4} + \frac{F^3}{27} \geq 0 ,$$

then

$$P + q^3 \geq 0 ,$$

which is satisfied when $q \geq 0$; i.e., when $a^2 r_0^2 + b^2 z_0^2 \geq (a^2 - b^2)^2$, which represents all space excluding the open region bounded by the ellipsoid

$$\frac{r_0^2}{a^2 e^4} + \frac{z_0^2}{b^2 e'^4} = 1.$$

Since $ae^2 < be'^2 < 43$ km, this region is of no practical interest. Therefore, $q \geq 0$ will be the constraint. Applying Theorem I to Equation (7) with $-t$ substituted for t , we see that Equation (7) has no negative roots. Hence, Equation (7) has only one real root and it is positive.

$$\begin{aligned} \sqrt{\frac{H^2}{4} + \frac{F^3}{27}} - \frac{H}{2} &= \frac{2}{6^3} \sqrt{P(P+q^3)} + \frac{1}{6^3} (2P+q^3) = \frac{1}{6^3} \{2P+q^3 + 2\sqrt{P(P+q^3)}\} \\ &= \frac{1}{6^3} (\sqrt{P+q^3} + \sqrt{P})^2. \end{aligned}$$

$$\therefore \left(\sqrt{\frac{H^2}{4} + \frac{F^3}{27}} - \frac{H}{2} \right)^{\frac{1}{3}} = \frac{Q}{6},$$

where

$$Q = (\sqrt{P+q^3} + \sqrt{P})^{\frac{2}{3}}.$$

Similarly,

$$\left(\sqrt{\frac{H^2}{4} + \frac{F^3}{27}} + \frac{H}{2} \right)^{\frac{1}{3}} = -\frac{Q'}{6},$$

where

$$Q' = (\sqrt{P+q^3} - \sqrt{P})^{\frac{2}{3}}.$$

(Note that $Q' = \frac{q^2}{Q}$. However, putting Q' in this form is undesirable as it introduces a singularity when $Q = 0$, which occurs when either both q and r_0 equal zero or both q and z_0 equal zero.)

Substituting these values into Equation (9),

$$t' = \frac{1}{6}(Q+Q') \Rightarrow t = \frac{1}{6}(q+Q+Q').$$

It will now be shown that $q \leq 2t \leq q + b^2$, where the left-hand equality occurs when either $r_0 = 0$ or $z_0 = 0$ and the right-hand equality occurs when $ar_0 = blz_0$.

Rewriting Equation (7) in the form

$$t^2(2t-q) = \frac{a^2 r_0^2 z_0^2}{e'^8},$$

the lower bound of t is deduced at once. Now let

$$g = \tau^3 - q\tau^2 - \frac{4a^2 r_0^2 z_0^2}{e'^8},$$

where $\tau \geq q$. Then

$$\frac{dg}{d\tau} = \tau(3\tau - 2q) \geq 0.$$

Thus, g is a monotonically increasing function of τ for all $\tau \geq q$. Since $2t \geq q$ and $g = 0$ when $\tau = 2t$, it follows that $g \geq 0$ when $\tau \geq 2t$. Putting $\tau = q + b^2$ in the expression for g ,

$$\begin{aligned} g|_{\tau=q+b^2} &= b^2(q + b^2)^2 - \frac{4a^2 r_0^2 z_0^2}{e'^8} = b^2 \left(p^2 + \frac{r_0^2}{e^2 e'^2} \right)^2 - \frac{4b^2 p^2 r_0^2}{e^2 e'^2}, \text{ using Equation (5)} \\ &= b^2 \left(p^2 - \frac{r_0^2}{e^2 e'^2} \right)^2 \geq 0 \end{aligned}$$

$\Rightarrow 2t \leq q + b^2$, equality occurring when $p = \frac{r_0}{ee'}$; i.e., when $\frac{r_0}{e} = \frac{|z_0|}{e'}$, or when $ar_0 = b|z_0|$, in which case,

$$q = 2p^2 - b^2, \quad t = \frac{1}{2}(q + b^2) = p^2.$$

Solving for c and d from the system of equations (6),

$$c = \sqrt{p^2 - q + 2t}, \quad d = \frac{p}{c}(t + b^2) = \sqrt{t^2 + p^2 b^2}. \quad (10)$$

Since $2t \geq q$, it follows that $c \geq p$. Now, $(|z|^2 + p|z| + t)^2 = (c|z| + d)^2$ implies $|z|^2 + p|z| + t = \pm (c|z| + d)$, from which the following two quadratics in $|z|$ are obtained:

$$|z|^2 + (p - c)|z| + t - d = 0, \quad (11)$$

and

$$|z|^2 + (p + c)|z| + t + d = 0. \quad (12)$$

It has already been shown that Equation (3) has exactly one positive root. Furthermore, since $t^2 - d^2 = -p^2 b^2 \leq 0$ and $d \geq 0$, it follows that $t - d \leq 0 \leq t + d$. Now, $t - d$ is the last term of Equation (11) and has just been shown to be negative. Hence, by applying Theorem II to Equation (11), we see immediately that Equation (11) must contain the required positive root. Hence,

$$|z| = \frac{-(p-c) + \sqrt{(p-c)^2 - 4(t-d)}}{2}$$

$$= \frac{1}{2} \left(c - p + \sqrt{2p^2 - q - 2t - 2pc + 4\sqrt{t^2 + p^2b^2}} \right).$$

This solution is free of singularities. It is easy to show that $\lim_{z_0 \rightarrow 0} |z| = 0$ and $\lim_{r_0 \rightarrow 0} |z| = b$. We have

$$\lim_{z_0 \rightarrow 0} p = 0, \quad \lim_{z_0 \rightarrow 0} P = \lim_{r_0 \rightarrow 0} P = 0 \Rightarrow \lim_{z_0 \rightarrow 0} Q = \lim_{z_0 \rightarrow 0} Q' = \lim_{r_0 \rightarrow 0} Q = \lim_{r_0 \rightarrow 0} Q' = q$$

$$\Rightarrow \lim_{z_0 \rightarrow 0} t = \lim_{r_0 \rightarrow 0} t = \frac{q}{2} \Rightarrow \lim_{z_0 \rightarrow 0} c = 0, \quad \lim_{r_0 \rightarrow 0} c = p.$$

$$\therefore \lim_{z_0 \rightarrow 0} |z| = \frac{1}{2} \sqrt{-2q + 2q} = 0,$$

$$\begin{aligned} \lim_{r_0 \rightarrow 0} |z| &= \frac{1}{2} \sqrt{2p^2 - 2q - 2p^2 + 4\sqrt{\frac{q^2}{4} + p^2b^2}} \\ &= \frac{1}{2} \sqrt{-2(p^2 - b^2) + 4\sqrt{\frac{1}{4}(p^2 - b^2)^2 + p^2b^2}} \\ &= \frac{1}{2} \sqrt{-2(p^2 - b^2) + 2(p^2 + b^2)} = b. \end{aligned}$$

Proceeding with the derivation,

$$r = a\sqrt{1 - \frac{z^2}{b^2}}, \quad x = x_0 \frac{r}{r_0}, \quad y = y_0 \frac{r}{r_0}, \quad \text{and} \quad \lambda = \tan^{-1} \frac{y_0}{x_0},$$

where $r_0 \neq 0$. (Here, λ is the longitude at (x,y) .)

To compute the geodetic latitude and altitude at $P'(x,y,z)$, it is desirable to introduce a geometric term, N_e , which is never zero. N_e is defined to be the distance along the ellipsoidal normal from the surface of the ellipsoid to the z -axis (see Figure 1).




FIGURE 1. ELLIPSOIDAL NORMAL

From Figure 1,

$$\cos \phi = \frac{r}{N_e}.$$

Also,

$$\tan \phi = \frac{a^2 z}{b^2 r}, \text{ from Equation (2)}$$

$$\begin{aligned} \Rightarrow N_e &= r \sec \phi = r \sqrt{1 + \tan^2 \phi} = r \sqrt{1 + \frac{a^4 z^2}{b^4 r^2}} = \sqrt{r^2 + \frac{a^4 z^2}{b^4}} \\ &= \sqrt{a^2 \left(1 - \frac{z^2}{b^2}\right) + \frac{a^4 z^2}{b^4}} = a \sqrt{1 + \frac{a^2 - b^2}{b^4} z^2} = a \sqrt{1 + \frac{e'^2 z^2}{b^2}}, \end{aligned} \quad (13)$$

and

$$\sin \phi = \cos \phi \tan \phi = \frac{a^2 z}{b^2 N_e} \Rightarrow \phi = \sin^{-1} \frac{a^2 z}{b^2 N_e},$$

which is the desired expression for the geodetic latitude.

Also from Figure 1,

$$h \cos \phi = r_0 - r = r_0 - N_e \cos \phi,$$

$$h \sin \phi = z_0 - z = z_0 - \frac{b^2}{a^2} N_e \sin \phi .$$

Multiplying the first equation by $\cos \phi$ and the second by $\sin \phi$ and adding,

$$h = r_0 \cos \phi + z_0 \sin \phi - N_e \left(\cos^2 \phi + \frac{b^2}{a^2} \sin^2 \phi \right) .$$

Now

$$\begin{aligned} \tan^2 \phi &= \frac{a^4 z^2}{b^4 r^2} = \frac{a^2 z^2}{b^2 (b^2 - z^2)} \\ \Rightarrow \cos^2 \phi + \frac{b^2}{a^2} \sin^2 \phi &= \frac{1 + \frac{b^2}{a^2} \tan^2 \phi}{1 + \tan^2 \phi} = \frac{1 + \frac{z^2}{b^2 - z^2}}{1 + \frac{a^2 z^2}{b^2 (b^2 - z^2)}} = \frac{b^4}{b^4 + (a^2 - b^2) z^2} \\ &= \frac{1}{1 + \frac{e'^2 z^2}{b^2}} = \frac{a^2}{N_e^2}, \text{ using Equation (13).} \end{aligned}$$

Hence

$$h = r_0 \cos \phi + z_0 \sin \phi - \frac{a^2}{N_e} ,$$

which is the desired expression for the geodetic altitude.

To recapitulate, given a , b , x_0 , y_0 , and z_0 , the algorithm is as follows:

$$e^2 = 1 - \frac{b^2}{a^2}, e'^2 = \frac{a^2}{b^2} - 1, r_0 = \sqrt{x_0^2 + y_0^2}, p = \frac{|z_0|}{e'^2}, q = p^2 - b^2 + \frac{r_0^2}{e^2 e'^2} .$$

If $q \geq 0$, then

$$P = \frac{27a^2 r_0^2 z_0^2}{e'^8}, Q = \left(\sqrt{P+q^3} + \sqrt{P} \right)^{\frac{2}{3}}, Q' = \left(\sqrt{P+q^3} - \sqrt{P} \right)^{\frac{2}{3}}, t = \frac{1}{6}(q + Q + Q'),$$

$$c = \sqrt{p^2 - q + 2t}, z = \frac{1}{2} \left(c - p + \sqrt{2p^2 - q - 2t - 2pc + 4\sqrt{t^2 + p^2 b^2}} \right) \text{sign } z_0 ,$$

$$N_e = a \sqrt{1 + \frac{e'^2 z^2}{b^2}}, \phi = \sin^{-1} \frac{a^2 z}{b^2 N_e}, h = r_0 \cos \phi + z_0 \sin \phi - \frac{a^2}{N_e} .$$

CONCLUSION

An efficient model for transforming Earth-centered, Earth-fixed coordinates to geodetic coordinates has been presented. The model systematically derives exact expressions for the geodetic latitude and altitude, which are free of singularities. To the author's best knowledge, no such expressions exist in the literature.

DISTRIBUTION

	<u>COPIES</u>		<u>COPIES</u>
DOD ACTIVITIES (CONUS)		INTERNAL	
ATTN SP231	1	E231	3
STRATEGIC SYSTEMS PROGRAMS		E282 SWANSBURG	1
DEPARTMENT OF THE NAVY		K10	3
WASHINGTON DC 20376-5002		K104 FELL	1
		K12 CARR	1
DEFENSE TECHNICAL INFORMATION		K12 O'TOOLE	1
CENTER		K13 EVANS	1
CAMERON STATION		K407 GATES	1
ALEXANDRIA VA 22304-6145	12	K41 SOFAIR	10
		K42	1
ATTN CODE E29L (TECHNICAL LIBRARY)	1	K43	1
COMMANDING OFFICER		K44	1
CSSDD NSWC		K52	1
6703 W HIGHWAY 98		N74 (GIDEP)	1
PANAMA CITY FL 32407-7001			
 NON-DOD ACTIVITIES (CONUS)			
ATTN GIFT AND EXCHANGE DIVISION	4		
LIBRARY OF CONGRESS			
WASHINGTON DC 20540			
 THE CNA CORPORATION			
PO BOX 16268			
ALEXANDRIA VA 22302-0268	1		
 ATTN RICHARD RAPP	1		
DEPARTMENT OF GEODETIC SCIENCE			
AND SURVEYING			
404 COCKINS HALL			
1958 NEIL AVE			
COLUMBUS OH 43210-1247			

REPORT DOCUMENTATION PAGE			Form Approved OBM No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, search existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE April 1995	3. REPORT TYPE AND DATES COVERED Final		
4. TITLE AND SUBTITLE Improved Method for Calculating Exact Geodetic Latitude and Altitude		5. FUNDING NUMBERS		
6. AUTHOR(s) Isaac Sofair				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commander Naval Surface Warfare Center, Dahlgren Division (Code K41) 17320 Dahlgren Road Dahlgren, VA 22448-5100		8. PERFORMING ORGANIZATION REPORT NUMBER NSWCDD/TR-94/77		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) An efficient model for transforming Earth-centered, Earth-fixed coordinates to geodetic coordinates is presented. The model systematically derives exact expressions for the geodetic latitude and altitude, which are free of singularities. To the author's best knowledge, no such expressions exist in the literature.				
14. SUBJECT TERMS Geodetic Latitude, Geodetic Altitude			15. NUMBER OF PAGES 18	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and its title page. Instructions for filling in each block of the form follow. It is important to **stay within the lines** to meet **optical scanning requirements**.

Block 1. Agency Use Only (Leave blank).

Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. *If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

Block 4. Title and Subtitle. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

Block 5. Funding Numbers. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit Accession No.

Block 6. Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

Block 7. Performing Organization Name(s) and address(es). Self-explanatory.

Block 8. Performing Organization Report Number. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.

Block 10. Sponsoring/Monitoring Agency Report Number. (If Known)

Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of ...; To be published in... . When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. Distribution/Availability Statement. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents"
DOE - See authorities.
NASA - See Handbook NHB 2200.2
NTIS - Leave blank

Block 12b. Distribution Code.

DOD - Leave blank.
DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.
NASA - Leave blank.
NTIS - Leave blank.

Block 13. Abstract. Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

Block 14. Subject Terms. Keywords or phrases identifying major subjects in the report.

Block 15. Number of Pages. Enter the total number of pages.

Block 16. Price Code. Enter appropriate price code (*NTIS only*).

Block 17-19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of this page.

Block 20. Limitation of Abstract. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited or SAR (same as report). An entry in this block is necessary if the abstract is to limited. If blank, the abstract is assumed to be unlimited.